

# Geminal wavefunctions: Looking for size extensivity beyond Slater-Jastrow

Pascal Bugnion

Cavendish Laboratory  
Cambridge University

# Quantum Monte-Carlo

- 1 Use VMC to generate a trial wavefunction.
- 2 Run a DMC calculation on this trial wavefunction in the **fixed-node** approximation.

# Accuracy

Improving the accuracy of DMC means improving the nodal surface of the trial wavefunction.

This is controlled by the **anti-symmetric part** of the wavefunction.

# Beyond Slater determinants?

- Backflow
- Multi-determinant expansion
- Geminal and Pfaffian wavefunctions

# Geminals

# Geminals

Idea

Can we build correlation into the orbitals?

# Geminals

## Idea

Can we build correlation into the orbitals?

This is where multi-electron orbitals come in. This talk will focus on orbitals containing two electrons of opposite spins, also called bi-orbitals or **geminals**.

$$\{\Phi_1(\mathbf{x}, \mathbf{y}), \Phi_2(\mathbf{x}, \mathbf{y}) \dots\}$$

# Geminal Parametrization





# Geminal Parametrization

Assume no spin-polarization, and each bi-orbital contains an up-spin electron  $\mathbf{x}$  and a down-spin electron  $\mathbf{y}$ .



$$\Phi(\mathbf{x}, \mathbf{y}) = \sum_{i,j} g_{ij} \cdot \phi_i(\mathbf{x}) \phi_j(\mathbf{y})$$

$$\Phi^\dagger = \sum_{i,j} g_{ij} \cdot b_i^\dagger \bar{b}_j^\dagger$$



# An Example: $H_4$





# An Example: $H_4$





# $H_4$ geminal basis states





# $H_4$ geminal basis states





# $H_4$ geminal basis states



# $H_4$ geminal basis states



# $H_4$ geminal basis states





# From bi-orbitals to wavefunctions

$$|\Phi\rangle = a \left( \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & - \end{array} \right\rangle + \left| \begin{array}{cc} - & - \\ - & \uparrow\downarrow \end{array} \right\rangle \right) \\ + b \left( \left| \begin{array}{cc} \uparrow\downarrow & - \\ - & - \end{array} \right\rangle + \left| \begin{array}{cc} - & \uparrow\downarrow \\ - & - \end{array} \right\rangle \right)$$

# From bi-orbitals to wavefunctions

$$|\Phi\rangle = a \left( \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & - \end{array} \right\rangle + \left| \begin{array}{cc} - & - \\ - & \uparrow\downarrow \end{array} \right\rangle \right) \\ + b \left( \left| \begin{array}{cc} \uparrow\downarrow & - \\ - & - \end{array} \right\rangle + \left| \begin{array}{cc} - & \uparrow\downarrow \\ - & - \end{array} \right\rangle \right)$$

$$\Phi^{\dagger 2} |0\rangle =$$

# From bi-orbitals to wavefunctions




$$|\Phi\rangle = a \left( \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & - \end{array} \right\rangle + \left| \begin{array}{cc} - & - \\ - & \uparrow\downarrow \end{array} \right\rangle \right) \\ + b \left( \left| \begin{array}{cc} \uparrow\downarrow & - \\ - & - \end{array} \right\rangle + \left| \begin{array}{cc} - & \uparrow\downarrow \\ - & - \end{array} \right\rangle \right)$$

$$\Phi^{\dagger 2} |0\rangle = a^2 \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & \uparrow\downarrow \end{array} \right\rangle + b^2 \left| \begin{array}{cc} \uparrow\downarrow & \uparrow\downarrow \\ - & - \end{array} \right\rangle + \dots$$




# From bi-orbitals to wavefunctions

$$\begin{aligned}\Phi^{\dagger 2} |0\rangle &= a^2 \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & \uparrow\downarrow \end{array} \right\rangle + b^2 \left| \begin{array}{cc} \uparrow\downarrow & \uparrow\downarrow \\ - & - \end{array} \right\rangle \\ &+ ab \cdot \left( \left| \begin{array}{cc} \uparrow\downarrow & - \\ - & \uparrow\downarrow \end{array} \right\rangle + \left| \begin{array}{cc} - & \uparrow\downarrow \\ \uparrow\downarrow & - \end{array} \right\rangle \right. \\ &\quad \left. + \left| \begin{array}{cc} \uparrow\downarrow & - \\ \uparrow\downarrow & - \end{array} \right\rangle + \left| \begin{array}{cc} - & \uparrow\downarrow \\ - & \uparrow\downarrow \end{array} \right\rangle \right)\end{aligned}$$

# Test system: $H_{2x}$




1.4a.u.




$b$

# H<sub>2x</sub> Results




# HEG Results

Homogeneous electron gas,  $r_s = 2 a_0$ .



# HEG Results


Homogeneous electron gas,  $r_s = 2 a_0$ .






# HEG Results: Geminal-Jastrow in DMC

Geminal-Jastrow vs. Slater-Jastrow:



# HEG Results: Geminal-Jastrow in DMC

Geminal-Jastrow vs. Slater-Jastrow:



# Multi-geminals?

Additional freedom can be introduced by introducing additional geminals.

# Multi-geminals?

Additional freedom can be introduced by introducing additional geminals.

For example, for a four electron wavefunction:

$$|\Psi\rangle = \Phi_1^{\dagger 2} |0\rangle + \Phi_2^{\dagger 2} |0\rangle + \dots$$

# Decoupling excitations

$$|\Phi_1\rangle = 0 \left| \begin{array}{cc} - & - \\ \uparrow\downarrow & - \end{array} \right\rangle + a \left| \begin{array}{cc} - & - \\ - & \uparrow\downarrow \end{array} \right\rangle \\ + b \left| \begin{array}{cc} \uparrow\downarrow & - \\ - & - \end{array} \right\rangle + c \left| \begin{array}{cc} - & \uparrow\downarrow \\ - & - \end{array} \right\rangle$$

# Decoupling excitations

$$|\Phi_1\rangle = a \begin{vmatrix} - & - \\ - & \uparrow\downarrow \end{vmatrix} + b \begin{vmatrix} \uparrow\downarrow & - \\ - & - \end{vmatrix} + c \begin{vmatrix} - & \uparrow\downarrow \\ - & - \end{vmatrix}$$

# Decoupling excitations

$$|\Phi_1\rangle = a \begin{vmatrix} - & - \\ - & \uparrow\downarrow \end{vmatrix} + b \begin{vmatrix} \uparrow\downarrow & - \\ - & - \end{vmatrix} + c \begin{vmatrix} - & \uparrow\downarrow \\ - & - \end{vmatrix}$$

$$\begin{aligned} \Phi_1^{\dagger 2} |0\rangle &= ab \begin{vmatrix} \uparrow\downarrow & - \\ - & \uparrow\downarrow \end{vmatrix} + ac \begin{vmatrix} - & \uparrow\downarrow \\ - & \uparrow\downarrow \end{vmatrix} \\ &\quad + bc \begin{vmatrix} \uparrow\downarrow & \uparrow\downarrow \\ - & - \end{vmatrix} \end{aligned}$$

# Multi-geminal wavefunction for $H_4$

$$|\Psi\rangle = |HF\rangle + \Phi_1^{\dagger 2} |0\rangle + \Phi_2^{\dagger 2} |0\rangle$$



# Multi-geminal wavefunction for $H_4$

$$\begin{aligned} |\Psi\rangle = & |HF\rangle + ab \cdot \left( \left| \begin{array}{c} \uparrow\downarrow - \\ - \uparrow\downarrow \end{array} \right\rangle + \left| \begin{array}{c} - \uparrow\downarrow \\ \uparrow\downarrow - \end{array} \right\rangle \right) \\ & + ac \cdot \left( \left| \begin{array}{c} - \uparrow\downarrow \\ - \uparrow\downarrow \end{array} \right\rangle + \left| \begin{array}{c} \uparrow\downarrow - \\ \uparrow\downarrow - \end{array} \right\rangle \right) \\ & + 2bc \left| \begin{array}{c} \uparrow\downarrow \uparrow\downarrow \\ - - \end{array} \right\rangle \end{aligned}$$

# Multi-geminal wavefunction

# Multi-geminal wavefunction

For a non-spin polarised system with  $2N$  electrons:

$$|\Psi\rangle = |HF\rangle + \sum_{\substack{a \in \text{up electrons} \\ b \in \text{down electrons}}} a_{ab} \cdot \Phi_{ab}^{\dagger N} |0\rangle$$

# Multi-geminal wavefunction

For a non-spin polarised system with  $2N$  electrons:

$$|\Psi\rangle = |HF\rangle + \sum_{\substack{a \in \text{up electrons} \\ b \in \text{down electrons}}} a_{ab} \cdot \Phi_{ab}^\dagger{}^N |0\rangle$$

where


$$\Phi_{ab}^\dagger = \sum_{i,j} g_{ij} \cdot b_i^\dagger \bar{b}_j^\dagger$$

where  $g_{ij} = 0$  if  $i = a$  or  $j = b$ .

# Multi-geminal results

# Multi-geminal results

Homogeneous electron gas,  $r_s = 2 a_0$ .



# Remaining questions

- 1 How good is the Multi-Geminal-Jastrow wavefunction in DMC?

# Remaining questions

- 1 How good is the Multi-Geminal-Jastrow wavefunction in DMC?
- 2 Is it any better than a multi-determinant expansion?



# Conclusions

- Single geminal wavefunctions give poor energies, both at VMC and DMC level.

# Conclusions

- Single geminal wavefunctions give poor energies, both at VMC and DMC level.
- The multi-geminal wavefunction presented here might be promising.

*Geminal wavefunctions with Jastrow correlation: a first application to atoms*

M. Casula and S. Sorella, J. Chem. Phys. 119, 6500 (2003)

*Generalized Pairing Wave Functions and Nodal Properties for Electronic Structure Quantum Monte Carlo*

Michael Bajdich (PhD thesis) - available online.

Thank you for your  
attention!